

Physics

Magnetic Field Dependence of $A \rightarrow B$ Phase Transition Temperature in Superfluid ${}^3\text{He}$

Giorgi Baramidze*, Giorgi Kharadze**

* E. Andronikashvili Institute of Physics, Tbilisi

** Academy Member, E. Andronikashvili Institute of Physics, Tbilisi

ABSTRACT. The magnetic field dependence of the $A \rightarrow B$ transition temperature T_{AB} in superfluid ${}^3\text{He}$ is considered in order to take into account the linear-in-field contribution. In the high field region, where the quadratic-in-field contribution prevails, the well-known answer is restored. On the other hand, it is pointed out that the Fermi liquid effects shift the observability of the linear-in-field region to rather low magnetic fields.

© 2009 Bull. Georg. Natl. Acad. Sci.

Key words: superfluidity of liquid ${}^3\text{He}$, magnetic field.

Soon after the discovery of the superfluidity of liquid ${}^3\text{He}$ it was established that near the critical temperature $T_C(H)$ the phase diagram of superfluid state undergoes a profound modification under the action of even small external magnetic fields [1,2,3]. This modification shows up in the elimination of the direct normal to B phase transition over the entire phase diagram in favour of an anisotropic A phase.

The $A \rightarrow B$ transition temperature $T_{AB} < T_C$ can be established by equating the free energies $F_A(H)$ and $F_B(H)$. Near the zero-field transition temperature $T_C(P)$ the free energy can be found by minimizing the Ginzburg-Landau functional

$$F_S = 3\alpha_{\mu\nu} \langle \Delta_\mu \Delta_\nu^* \rangle + F_S^{(4)}(\bar{\Delta}), \quad (1)$$

where the fourth order contribution in the order-parameter $\bar{\Delta}(\hat{k})$ reads as

$$F_S^{(4)} = 9 \left\{ \beta_1 \langle \Delta_\mu \Delta_\mu \rangle \langle \Delta_\nu^* \Delta_\nu^* \rangle + \beta_2 \langle |\bar{\Delta}|^2 \rangle^2 + \beta_3 \langle \Delta_\mu \Delta_\nu \rangle \langle \Delta_\mu^* \Delta_\nu^* \rangle + \beta_4 \langle \Delta_\mu \Delta_\nu^* \rangle \langle \Delta_\mu^* \Delta_\nu \rangle + (2) \right. \\ \left. + \beta_5 \langle \Delta_\mu \Delta_\nu^* \rangle \langle \Delta_\mu \Delta_\nu^* \rangle \right\}.$$

In Eqs. (1) and (2) the spin vector $\bar{\Delta}$ is defined according to the general expression for the order-parameter $A_{\mu i}$ of the spin triplet P -wave Cooper condensate: $\Delta_\mu(\hat{k}) = A_{\mu i} \hat{k}_i$, and the angular brackets $\langle \dots \rangle$ stand for an average over the momentum direction \hat{k} on the Fermi surface. In the presence of a magnetic field \vec{H} the tensor coefficient $\alpha_{\mu\nu}$ of the second order term in Eq.(1) reads as:

$$\alpha_{\mu\nu} = \alpha \delta_{\mu\nu} + i g_1 \varepsilon_{\mu\nu\lambda} H_\lambda + g_2 H_\mu H_\nu, \quad (3)$$

where:

$$\alpha = \frac{1}{3} N_F \ln \left(\frac{T}{T_C} \right), \quad (4a)$$

$$g_1 H = -\frac{1}{3} N_F \eta h, \quad (4b)$$

$$g_2 H^2 = \frac{1}{3} N_F \kappa h^2, \quad (4c)$$

with $h = \hbar \gamma H / (2k_B T_C) = H / H_0$. The dimensionless coefficients η and κ could be considered as phenomeno-

logical quantities although their values can be estimated according to microscopic calculations. In the weak-coupling approximation [4]

$$\eta_{wc} = \frac{N'_F}{N_F} k_B T_C \ln \left(\frac{2\gamma_E \hbar \omega_C}{\pi k_B T_C} \right), \quad (5)$$

where N'_F stands for the derivative of the quasiparticle DOS with respect to the energy at the Fermi level. Detailed calculations which take into account the linear-in-field corrections to the Fermi liquid parameters are performed in Ref.[5].

As to the κ coefficient, it stems from the free energy part

$$\delta F_H^{(2)} = \frac{1}{2} \delta \chi_s H^2, \quad \delta \chi_s = \chi_s - \chi_N, \quad (6)$$

where χ_s (χ_N) stands for the magnetic susceptibility of the superfluid (normal) phase. For the B phase near T_C ,

$\delta F_H^{(2)} = g_2 H^2 \Delta^2$ where $g_2 H^2$ is given according to Eq. (4c) with

$$\kappa = \frac{7\zeta(3)}{4\pi^2} \frac{1}{(1 + F_0^a)^2}. \quad (7)$$

Here the Fermi liquid parameter $F_0^a \cong -3/4$ and is weakly pressure dependent.

As is well known, the linear-in-field term (4b) is the origin of a tiny splitting of the A phase (due to a small asymmetry of the density of quasiparticle states at the Fermi level). The quadratic-in-field contribution (4c) gives rise to the magnetic field suppression of the $\Delta_{\uparrow\downarrow}$ component of the energy gap of the B phase.

Based on the argument that the term (4b) is rather small, in the majority of considerations of $T_{AB} = T_{AB}(H)$ this term is usually discarded, and in a standard way one starts from the following expressions for the equilibrium free energies of the A and B phases:

$$F_A = -\frac{1}{4\beta_{245}} \alpha^2, \quad (8a)$$

$$F_B = -\frac{1}{2(3\beta_{12} + \beta_{345})} \times \left[\frac{3}{2} \alpha^2 + g_2 H^2 \alpha + \frac{2\beta_{12} + \beta_{345}}{2\beta_{345}} (g_2 H^2)^2 \right]. \quad (8b)$$

Since here the linear-in-field contribution (proportional to g_1) is dropped, the action of the magnetic field appears only in F_B .

Equating (8a) and (8b), it is found that $T_{AB}(H)$ is to be obtained from the equation:

$$\varphi = \alpha^2 + a\alpha + b = 0 \quad (9)$$

with

$$a = 2P_1 g_2 H^2, \quad (10)$$

$$b = P_1^2 (1 - q_1) (g_2 H^2)^2,$$

where the coefficients P_1 and q_1 are defined in the Appendix. From Eq. (9) follows the answer for $\tau_{AB} = 1 - T_{AB}/T_C$:

$$\tau_{AB} = P_1 (1 + \sqrt{q_1}) \kappa h^2 \quad (11)$$

which reproduces a well-known result obtained in Ref.[6].

Now we turn to the role of the linear-in-field contribution to $T_{AB}(H)$. This question was first posed in Ref. [7]. The starting point is the construction of the expressions for $F_{A2}(H)$ and $F_B(H)$, the contribution (4b) being taken into account. In a standard way it is established that

$$F_{A2} = -\frac{1}{4\beta_{245}} \left[\alpha^2 - \frac{\beta_{245}}{\beta_5} g_1^2 H^2 \right], \quad (12)$$

$$F_B = -\frac{1}{2(3\beta_{12} + \beta_{345})} \times \left\{ \frac{3}{2} \alpha^2 + g_2 H^2 \alpha + \frac{3\beta_{12} + \beta_{345}}{\beta_4 - (3\beta_1 + \beta_{35})} g_1^2 H^2 + \frac{2\beta_{12} + \beta_{345}}{2\beta_{345}} (g_2 H^2)^2 + \frac{(3\beta_{12} + \beta_{345})^2}{(3\beta_1 + \beta_{35} - \beta_4)^2} \times \left[\frac{\beta_1}{\beta_{345}} \frac{g_1^2 g_2 H^4}{\alpha} + \frac{\beta_1 (3\beta_{12} + \beta_{345})}{4(3\beta_1 + \beta_{35} - \beta_4)^2} \frac{g_1^4 H^4}{\alpha^2} \right] \right\}. \quad (13)$$

Comparison of $F_{A2}(H)$ and $F_B(H)$ gives an equation for $T_{AB}(H)$

$$\varphi \left(\frac{T_{AB}}{T_C} \right) = \alpha^4 + a\alpha^3 + b\alpha^2 + c\alpha + d = 0, \quad (14)$$

where now

$$a = 2P_1 g_2 H^2,$$

$$b = -P_1 P_2 P_3 g_1^2 H^2 + P_1^2 (1 - q_1) (g_2 H^2)^2, \quad (15)$$

$$c = -P_1 P_2^2 P_4 g_1^2 g_2 H^4,$$

$$d = -P_1 P_2^3 P_5 (g_1 H^4).$$

The definition of all coefficients P_a is given in the Appendix.

In Ref.[7] the second and third lines in Eq.(13), containing terms on the order H^4 , were neglected. This has the following influence on the answer for $T_{AB}(H)$: i) neglecting of the contribution proportional to $(g_2 H^2)^2$

makes it impossible to reproduce the correct answer given by Eq. (11) (the term $\sqrt{q_1}$ will be lost), ii) neglecting of the contribution collected in the square brackets of the third line of Eq.(13) changes the answer for the linear-in-field contribution to $T_{AB}(H)$. To avoid these drawbacks we address Eq.(14) and, as a first step, perform the variable transformation $\alpha \rightarrow x - \frac{1}{4}a$, after which an equation for $x(T_{AB}/T_C)$ is obtained:

$$\varphi\left(\frac{T_{AB}}{T_C}\right) = x^4 + Ax^2 + Bx + C = 0 \quad (16)$$

with the coefficients

$$\begin{aligned} A &= -\left[P_1 P_2 P_3 (g_1 H)^2 + \frac{1}{2} P_1^2 (1 + 2q_1) (g_2 H^2)^2 \right], \\ B &= q_1 P_1^3 (g_2 H^2)^3 + q_2 P_1 P_2 g_1^2 g_2 H^4, \\ C &= -P_1 P_2^3 P_3 (g_1 H)^4 + \frac{1}{16} P_1^4 (1 - 4q_1) (g_2 H^2)^4 + \\ &\quad + \frac{1}{4} P_1^2 P_2 (P_1 P_3 - 2q_2) (g_1 g_2 H^3)^2. \end{aligned} \quad (17)$$

It is to be noted that as a result of the variable transformation used, in Eq.(16) the cubic term is absent. At the same time the coefficient of the linear term B is zero in the weak-coupling approximation.

In order to solve Eq.(16) we use the decomposition $x = x_0 - \sqrt{q_1} P_1 g_2 H^2$ which, generates the decomposition $\varphi = \varphi_0 + \delta\varphi$ with φ_0 being the solution of the equation

$$\varphi_0 = x_0^4 + A_0 x_0^2 + C_0 = 0, \quad (18)$$

where A_0 and C_0 are the coefficients A and C taken at $q_1 = q_2 = 0$

From Eq. (18) it is found that

$$x_0\left(\frac{T_{AB}}{T_C}\right) = -\sqrt{P_0^2 (g_1 H)^2 + \frac{1}{4} P_1^2 (g_2 H^2)^2}, \quad (19)$$

where

$$P_0^2 = \frac{1}{2} P_1 P_2 P_3 \left[1 + \sqrt{1 + 4 \frac{P_2 P_3}{P_1 P_3^2}} \right]. \quad (20)$$

Direct inspection shows that $\delta\varphi$ is the sum of terms proportional to the powers of q and the minimal power in H contained in $\delta\varphi$ is H^5 . For this reason we can use an approximation with $\delta\varphi$ disregarded, and as a result it is found that




Fig. The bold curve corresponds to the Landau parameter

$$F_0^a = -3/4 \text{ and the dashed curve is constructed at } F_0^a = 0.$$

$$\begin{aligned} \alpha\left(\frac{T_{AB}}{T_C}\right) &= x - \frac{1}{4}a = x_0 - \sqrt{q_1} P_1 g_2 H^2 - \frac{1}{2} P_1 g_2 H^2 = \\ &= -P_1 \left(\frac{1}{2} + \sqrt{q_1} \right) g_2 H^2 - \sqrt{P_0^2 (g_1 H)^2 + \frac{1}{4} P_1^2 (g_2 H^2)^2}. \end{aligned} \quad (21)$$

Using this result, we finally have the following simple answer for τ_{AB} :

$$\tau_{AB} = P_1 \left(\frac{1}{2} + \sqrt{q_1} \right) \kappa h^2 + \sqrt{P_0^2 (\eta h)^2 + \frac{1}{4} P_1^2 (\kappa h^2)^2}. \quad (22)$$

By introducing a characteristic magnetic field $H_* = 2(P_0/P_1)(\eta/\kappa)H_0$ the asymptotic regions where the quadratic-in-field ($H \gg H_*$) and the linear-in-field ($H \ll H_*$) contributions to T_{AB} prevail are found:

$$\tau_{AB} = \begin{cases} P_1 (1 + \sqrt{q_1}) \kappa h^2, & H \gg H_* \\ P_0 \eta h, & H \ll H_* \end{cases} \quad (23)$$

For $H \gg H_*$ the well-known result [6,8] is reproduced.

In order to isolate the role of the linear-in-field contribution to T_{AB} it is convenient to consider

$$\delta\tau_{AB}(h) = \tau_{AB}(h) - (1 + \sqrt{q_1}) P_1 \kappa h^2, \quad (24)$$

and construct graphically

$$f(h) = \frac{\delta\tau_{AB}(h)}{h} = P_0 \eta \left[\sqrt{1 + \left(\frac{h}{h_*} \right)^2} - \frac{h}{h_*} \right], \quad (25)$$

where the scaling value $h_* = 2(P_0/P_1)(\eta/\kappa)$.

In the Figure $f(h)$ is plotted for the pressure

$P = 10 \text{ bar}$ ($P_0 \eta \approx 5,4 \cdot 10^{-2}$, $h_* \approx 1,6 \cdot 10^{-2}$). It is evident that the Fermi liquid effect shifts the linear-in field contribution to rather small values of $H \ll H_* = h_* H_0 \approx 420 \text{ G}$.

Appendix

The coefficients P_a ($a = 1, 2, 3, 4, 5$) and q_a ($a = 1, 2$), introduced in the main text, are defined as follows:

$$P_1 = \frac{\beta_{245}}{2\beta_{345} - 3\beta_{13}}, \quad P_2 = \frac{3\beta_{12} + \beta_{345}}{\beta_4 - 3\beta_1 - \beta_{35}},$$

$$P_3 = \frac{\beta_{45} - 3\beta_1 - \beta_3}{-\beta_5}, \quad P_4 = \frac{-2\beta_1}{\beta_{345}},$$

$$P_5 = \frac{-\beta_1}{2(\beta_4 - 3\beta_1 - \beta_{35})}.$$

$$q_1 = \frac{(3\beta_{12} + \beta_{345})(2\beta_{13} - \beta_{345})}{\beta_{245}\beta_{345}}, \quad q_2 = P_1 P_3 - P_2 P_4,$$

where $\beta_{ij\dots} = \beta_i + \beta_j + \dots$

The coefficients q_1 and q_2 contain only the strong-coupling corrections $\delta\beta_i$ defined according to the decomposition

$$\beta_1 = -\beta_0 + \delta\beta_1, \quad \beta_2 = 2\beta_0 + \delta\beta_2, \quad \beta_3 = 2\beta_0 + \delta\beta_3,$$

$$\beta_4 = 2\beta_0 + \delta\beta_4, \quad \beta_5 = -2\beta_0 + \delta\beta_5,$$

$$\beta_0 = \frac{7\zeta(3)}{240} \frac{N_F}{(\pi k_B T_C)^2}.$$

Acknowledgements

The authors highly appreciate the valuable discussions with the late Dr. Shota Nikolaishvili.

ფიზიკა

ზედენად ${}^3\text{He}$ -ში $A \rightarrow B$ ფაზური გადასვლის ტემპერატურის დამოკიდებულება მაგნიტურ ველზე

გ. ბარამიძე*, გ. ხარაძე**

* ე. ანდრონიკაშვილის ფიზიკის ინსტიტუტი, თბილისი

** აკადემიკოსი, ე. ანდრონიკაშვილის ფიზიკის ინსტიტუტი, თბილისი

განხილულია ზედენად ${}^3\text{He}$ -ში $A \rightarrow B$ ფაზური გადასვლის T_{AB} ტემპერატურის მაგნიტურ ველზე დამოკიდებულების თავისებურებანი. ნაჩვენებია, რომ ხსენებული დამოკიდებულების წრფივი მდგენელის ექსპერიმენტული დაშვება შესაძლებელია მხოლოდ საკმარისად მცირე მაგნიტური ველების პირობებში.

REFERENCES

1. *W.J. Gully et al.* (1973), *Phys. Rev. A* 8: 1633.
2. *D.N. Paulson et al.* (1974), *Phys. Rev. Lett.* 32: 1098.
3. *J.D. Feder et al.* (1981), *Phys. Rev. Lett.* 47: 428.
4. *V. Ambegaokar and D. Mermin* (1973), *Phys. Rev. Lett.* 30: 81.
5. *K. Bedell and K. Quadar* (1984), *Phys. Rev. B* 30: 2894.
6. *A.L. Fetter* (1975), In: *Quantum Statistics and the Many-Body Problem*, Plenum, NY; (1976) *J. Low Temp. Phys.* 23, 245.
7. *K. Levin and O. Walls* (1977), *Phys. Rev. B* 15: 4256.
8. *Y.H. Tang et al.* (1991), *Phys. Rev. Lett.* 67: 1775.

Received August, 2009